

Identification of known leaf rust resistance genes in bread wheat cultivars from China

XIAOCUI YAN¹, TAKELE-WELDU GEBREWAHID², RUI DONG¹, XING LI¹,
PEIPEI ZHANG¹, ZHANJUN YAO^{3*}, ZAIFENG LI^{1*}

¹Department of Plant Pathology, College of Plant Protection, Hebei Agricultural University, Baoding, Hebei, P.R. China

²College of Agriculture, Aksum University, Shire-Indaslassie, Tigray, Ethiopia

³College of Agronomy, Hebei Agricultural University, North China Key Laboratory for Crop Germplasm Resources of China's Education Ministry, Baoding, Hebei, P.R. China

*Corresponding authors: yzhj201@aliyun.com; lz7551@aliyun.com

Citation: Yan X., Gebrewahid T.-W., Dong R., Li X., Zhang P., Yao Z., Li Z. (2021): Identification of known leaf rust resistance genes in bread wheat cultivars from China. Czech J. Genet. Plant Breed., 57: 91–101.

Abstract: Leaf rust caused by *Puccinia triticina* Eriks. (*Pt*) is one of the most devastating fungal pathogens affecting wheat (*Triticum aestivum* L.) production worldwide. Deployment of resistant cultivars is the most environmentally friendly approach to control the disease. In this study, thirty-seven wheat lines from the Hubei and Shaanxi provinces in China were evaluated for seedling resistance in the greenhouse using eighteen *Pt* races. These lines were also tested for slow rusting resistance in the field in the 2014 to 2018 growing seasons. Eleven molecular markers closely associated with known *Lr* genes were used as part of the postulation process. Seven known *Lr* genes, 1, 13, 18, 14a, 26, 34 and 46 either singly or in combination were postulated in twenty-five cultivars. *Lr1* and *Lr26* were the most commonly identified genes detected in thirteen and ten cultivars, respectively. *Lr13* and *Lr46* were each found in four and five cultivars. *Lr34* was present in three cultivars. *Lr18* and *Lr14a* were identified in cultivar Xi'nong 538. Six cultivars displayed slow rusting resistance in the field tests. The resistant cultivars identified in the present study can be used as resistance parents in crosses aimed at pyramiding and the deployment of leaf rust resistance genes in China.

Keywords: adult plant resistance; gene postulation; molecular markers; *Triticum aestivum*

Bread wheat provides about 20% of the calories consumed by humankind (Fu et al. 2009). Wheat leaf rust caused by *Puccinia triticina* Eriks. (*Pt*), is one of the most important wheat diseases in many regions worldwide. This disease occurs in almost all wheat-growing areas, including North America, Europe, Asia, Australia, etc. (Dehne & Oerke 1998), and causes severe yield losses ranging from 30 to 50% (McIntosh et al. 1995). A particularly severe leaf rust epidemic in north-western Mexico caused an estimated yield loss of up to 70% during the 1970s (Dubin & Torres 1981). Widespread damaging leaf rust epidemics in China

were recorded in 1969, 1973, 1975 and 1979 (Dong 2001; Li et al. 2014). In 2012 and 2015, leaf rust caused destructive yield losses in the major wheat production regions of China, especially in North China (Zhou et al. 2013; Peng et al. 2016). Deployment of resistant cultivars is an economical and environmentally friendly way to control wheat leaf rust. To ensure the continuing effectiveness, it is important to identify and utilise new sources of leaf rust resistance (*Lr*) genes in wheat breeding programmes.

The principle of gene postulation is based on the gene-for-gene concept proposed by Flor (1955) to

identify resistance genes to the disease possibly present in the tested cultivars. A specific resistance gene in a host cultivar can be postulated from the response array produced by a series pathogen races with known avirulence/virulence characteristics. In addition, many resistance genes can be postulated using molecular markers although not all markers are fully diagnostic of the gene to which they are linked. *Lr* genes with reliable markers include 1, 10, 9, 19, 20, 24, 34, and 46. Numerous researchers have used these multi-race tests to postulate the *Lr* genes in different sets of wheat cultivars. For example, Yuan et al. (1992) identified the *Lr* genes 1, 3, 3bg, 9, 10, 13, 16, 23, 26, and 34 in forty-seven wheat cultivars using seventeen *Pt* races. Gebrewahid et al. (2017) identified twelve *Lr* genes, 1, 26, 3ka, 11, 10, 2b, 13, 21, 34, 37, 44, and 46 in eighty-three Chinese common wheat cultivars using eighteen *Pt* races. Wu et al. (2019) postulated six *Lr* genes (1, 26, 33, 34, 45 and 46) in forty-four wheat accessions using twenty *Pt* races.

Currently, there is limited information regarding leaf rust resistance genes in the leading contemporary cultivars grown in some regions of China. In this study, thirty-seven wheat cultivars grown in the Hubei and Shaanxi provinces were subjected to multi-race seedling tests, field tests and molecular marker detection. The objective of this study was to identify *Lr* genes in thirty-seven wheat cultivars from China.

MATERIAL AND METHODS

Plant materials and *Pt* races. Thirty-seven wheat cultivars from the Shaanxi and Hubei were used in this study. The regions and pedigrees of the thirty-seven cultivars are listed in Table 1. All the cultivars were tested for seedling response using eighteen *Pt* races in the greenhouse (Tables 2 and 3), and for slow rusting to leaf rust in the field during the 2014 to 2018 growing seasons. The International Maize and Wheat Improvement Center (CIMMYT) line, Saar with typical slow leaf rusting (Lillemo et al. 2008; Zhuang et al. 2009) and the highly susceptible line, Zhengzhou 5389 (final disease severity (FDS) > 90%) were used as the slow rusting and susceptible controls, respectively. Thirty-six lines with known *Lr* genes were utilised as a reference base to compare the seedling infection types (ITs) produced by each test line (Table 2). All the *Pt* races were named following the three-letter coding system of Long and Kolmer

(1989), with the addition of a fourth letter for the fourth set of test differentials (http://www.ars.usda.gov/SP2/UserFiles/ad_hoc/36400500/Cereal_rusts/pt_nomen.pdf). The seeds of all the thirty-seven test cultivars, thirty-six differential lines with known leaf rust resistance genes, susceptible line Zhengzhou 5389, and CIMMYT line Saar were provided by the Wheat Rust Laboratory of Hebei Agricultural University.

Seedling test. All the cultivars were planted in a greenhouse and inoculated with eighteen *Pt* races for the gene postulation (Tables 2 and 3). The gene postulation was conducted following the method described by Singh et al. (1999) with minor modifications. The seedling inoculations were performed by brushing urediniospores from sporulating susceptible seedlings onto the test seedlings when the first leaves were fully expanded. The inoculated seedlings were placed in plastic-covered cages and incubated at 18 °C and 100% relative humidity (RH) for 24 h before being transferred to a growth chamber maintained with 12 h light/12 h darkness at 18 °C to 20 °C with 70% RH. The ITs were scored 10 to 14 days post-inoculation according to the 0 to 4 infection type scale as modified by Roelfs et al. (1992).

Adult plant tests. All the thirty-seven cultivars, along with the susceptible control, Zhengzhou 5389 and the slow rusting check, Saar were planted in a randomised complete block design with two replicates at Zhoukou, in Henan Province and Baoding, in Hebei Province during the 2014 to 2018 cropping seasons. Approximately fifty seeds of each line were sown in 1.5 m single-row plots with 0.3-m spacing. Spreader rows of Zhengzhou 5389 were planted perpendicular and adjacent to the test rows. The field inoculation was conducted using a mixture of an equal amount of urediniospores from the FHRT, THTT and THJT *Pt* races suspended in 0.03% Tween 20 onto the spreader rows at the tillering stage. The disease severities as a percentage of the leaf area covered with uredinia were scored three times at about the 1-week interval with the first scoring 4 weeks after inoculation (Feekes growth stage 10) in each environment according to the modified Cobb scale (Peterson et al. 1948). The FDS was collected when the susceptible control, Zhengzhou 5389 was fully infected. The FDS data evaluation to leaf rust response was conducted following the methods described by Li et al. (2010).

Statistical analysis. The analysis of variance (ANOVA) and for determining the least significance

<https://doi.org/10.17221/6/2021-CJGPB>

differences (LSDs) for comparing the FDS values among the cultivars were performed using the IBM SPSS Statistics 19.0 software. Cultivars susceptible to the mixed *Pt* races at the seedling stage and having FDS values significantly lower than the slow rusting control, Saar in the field trials were considered to be slow rusting cultivars.

Molecular makers testing. The genomic DNA was extracted following the cetyltrimethyl-ammonium bromide (CTAB) method (Sharp et al. 1988). Eleven molecular markers reported as diagnostic or closely linked to nine *Lr* genes 1, 9, 10, 19, 20, 24, 26, 34

and 46 were used to test all the cultivars (Table 3). The polymerase chain reaction (PCR) assay was conducted following the protocol of Helguera et al. (2000), in a 20 µL reaction volume containing: 10 µL of 2 × TaqPCR Master Mix (Tiangen Biochemical Incorporation, Beijing), 6 µL of ddH₂O, 2 µL (4 mol per µL) of the primer, and 2 µL 4 ng/µL of the template DNA. All PCR amplification conditions are listed in Table 4. The amplified products were detected by 1.5% agarose gel electrophoresis or 12% non-denaturing polyacrylamide gel electrophoresis in the case of *Lr46*.

Table 1. The region and pedigree of 37 Chinese wheat cultivars tested for the leaf rust response

| Line No. | Genotype | Region | Pedigree | Line No. | Genotype | Origin | Pedigree |
|----------|---------------|---------|--|----------|---------------|---------|--|
| 1 | E'mai 580 | Hubei | (Taigu sterile lines)/ (wheat line 957565) | 20 | Shaanmai 139 | Shaanxi | Xiaoyan 22 × 94156/ N9134 F1 |
| 2 | E'mai 17 | Hubei | E'mai 12 variation plant | 21 | Xi'nong 538 | Shaanxi | Lankao 90(6)52-30/ Xiaoyan 6/Huaihe 9412 |
| 3 | E'mai 12 | Hubei | 750025-12/E'mai 6 | 22 | Xiaoyan 166 | Shaanxi | 87135-1-3-2-1-2/88111 |
| 4 | E'mai 18 | Hubei | SKUA/865146//E'mai 11 | 23 | Shaannong 138 | Shaanxi | Xinmai 9/HangShaan 354 |
| 5 | Xi'nong 291 | Shaanxi | (Xiaoyan 5) 4DN/ hexaploid triticales WOH45 | 24 | Xiaoyan 54 | Shaanxi | Xiaoyan 6 |
| 6 | Xiaoyan 22 | Shaanxi | (Xiaoyan 6/775-1)/ Xiaoyan 107 | 25 | Shaan 253 | Shaanxi | Shaan 229/Shaan 213 |
| 7 | Shaan 150 | Shaanxi | (4/6811(2)F7/8435-1-1-8) F1/Xiaoyan 6 | 26 | Xi'nong 6028 | Shaanxi | Jingyan 60/Zhongnong 28 |
| 8 | Gaoyou 503 | Shaanxi | 78506/84s504 | 27 | Shaanmai 159 | Shaanxi | Xiaoyan 597/89605 |
| 9 | Xi'nong 88 | Shaanxi | <i>Aegilops variabilis</i> cytoplasm | 28 | Xiaoyan 216 | Shaanxi | Lankao 906/ Xiaoyan 22 |
| 10 | Xi'an 93991 | Shaanxi | NA | 29 | Xi'nong 2611 | Shaanxi | Shaan229/ [84(14)43/83(2)3-3]/ (Xinong65 × Xiaoyan) |
| 11 | E'mai16 | Shaanxi | The variation of 7023 plant | 30 | Shaan 558 | Shaanxi | Xiaoyan 22/v9511 |
| 12 | Xi'nong 2000 | Shaanxi | Xi' nong 2611/386/ Xiaoyan22/Shaan 354 | 31 | Xi'nong 9871 | Shaanxi | Xinong 2208/Xiaoyan 22 |
| 13 | Xiaoyan 4 | Shaanxi | Fengchang 1/Xiaoyan 759 | 32 | Xi'nong 889 | Shaanxi | E Han-4/(Xiaoyan 6/ Xiaoyan 83352) |
| 14 | Shaan 627 | Shaanxi | NA | 33 | Xiaoyan 228 | Shaanxi | NA |
| 15 | Xi'nong 223 | Shaanxi | Xi'nong 389 variation plant | 34 | Xiaoyan 319 | Shaanxi | NA |
| 16 | Xi'nong 126 | Shaanxi | NA | 35 | Shaan 160 | Shaanxi | Shaan 213/winter wheat lines 167-6-4 |
| 17 | Shaannong 981 | Shaanxi | NA | 36 | Shaanmai 175 | Shaanxi | NA |
| 18 | Xi'nong 3517 | Shaanxi | Xi'nong 1376/Xi'nong 88 | 37 | Shaan 512 | Shaanxi | Shaanmai 150(A2) |
| 19 | Shaanken 6 | Shaanxi | Lankao 906/Xiaoyan 22 | | | | |

NA – not available

<https://doi.org/10.17221/6/2021-CJGPB>Table 2. Infection types produced by 36 reference lines with the single leaf rust resistance genes inoculated with 18 *Puccinia triticina* (*Pt*) races

| No. | <i>Lr</i> gene(s) | Infection types ^a to <i>Pt</i> races | | | | | | | | | | | | | | | | | |
|-----|----------------------|---|------|----|------|-----|------|------|-----|-----|------|------|------|-----|------|------|----|----|------|
| | | PH | TH | PH | KH | PH | THTT | KH | FH | FH | PHTT | THTT | PHTT | FH | FHHT | FHHT | TG | FH | FG |
| | | GQ | JT | JT | JS | JS | I | HT | RT | JQ | I | II | II | TR | I | II | GT | TT | MT |
| 1 | <i>TcLr1</i> | 4 | 3+ | 3+ | ;1 | 4 | 4 | 1 | 1 | 1 | 4 | 4 | 4 | 1 | 1 | 1 | 4 | 1 | 1 |
| 2 | <i>TcLr2a</i> | 1 | 3+ | 2+ | 3 | 1 | 4 | 4 | 1 | 1 | 1 | 4 | 1 | 0,; | 1 | 1 | 4 | ; | 1 |
| 3 | <i>TcLr2C</i> | 4 | 4 | 4 | 3+ | 4 | 4 | 3+ | 4 | 4 | 4 | 4 | 4 | 3+ | 4 | 4 | 3+ | 3+ | 4 |
| 4 | <i>TcLr3</i> | 4 | 4 | 4 | 3+ | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 3 | 4 | 4 |
| 5 | <i>TcLr9</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 6 | <i>TcLr16</i> | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 3+ | 4 | 3 | 3 | 3 | 3 | 3 | 3+ | 3+ |
| 7 | <i>TcLr24</i> | ; | ; | ; | ; | ; | ; | ; | ; | ; | ; | ; | ; | ; | ; | ; | ; | 0 | 0 |
| 8 | <i>TcLr26</i> | 3+ | 3+ | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 3 | 3 | 3 | 3 | 3 | 3+ | 2 | 3+ | 2 |
| 9 | <i>TcLr3ka</i> | 1 | 1 | 1 | 2 | ;;1 | 3 | 1 | 4 | 1 | 3 | 3 | 3c | 3+ | 2 | 2 | 1 | 3 | 3 |
| 10 | <i>TcLr11</i> | 4 | 4 | 3+ | 3 | 3 | 4 | 3 | 4 | 4 | 4 | 4 | 3c | 4 | 3+ | 3 | 3+ | 3+ | 2+ |
| 11 | <i>TcLr17</i> | 2 | 3c | 2+ | 1 | 3 | 4 | 1 | 1 | 1 | 3+ | 4 | 3 | 3 | 2+ | 1 | 2+ | 3 | 2 |
| 12 | <i>TcLr30</i> | 1 | 1 | 2+ | 1 | 1 | 3+ | 3c | 3 | 1 | 3+ | 4 | 4 | 4 | 3+ | 4 | 1 | 3 | 4 |
| 13 | <i>TcLrB</i> | 4 | 3+ | 4 | 4 | 4 | 4 | 3+ | 4 | 4 | 4 | 4 | 3 | 4 | 4 | 3+ | 4 | 3+ | 4 |
| 14 | <i>TcLr10</i> | 3+ | 4 | 4 | 4 | 4 | 4 | 3+ | 4 | 4 | 4 | 3c | 4 | 3 | 3 | 3 | 4 | 4 | 4 |
| 15 | <i>TcLr14a</i> | X | 4 | 3+ | 4 | 3+ | 4 | 4 | 3+ | X | 4 | 3+ | 4 | X | 3+ | 3 | 3+ | 3+ | 3+ |
| 16 | <i>TcLr18</i> | 1 | 3+ | 3 | 1 | 2 | 3 | 3+ | 3 | 2 | 4 | 4 | 4 | 3 | 4 | 4 | 3+ | 4 | 3+ |
| 17 | <i>TcLr2b</i> | 2 | 4 | 3+ | 4 | 3+ | 3+ | 3+ | 2+ | 3 | 3+ | 4 | 3 | 3+ | 3+ | 3 | 3+ | 4 | 3+ |
| 18 | <i>TcLr3bg</i> | 4 | 4 | 3+ | 4 | 4 | 4 | 3+ | 3+ | 3+ | 4 | 4 | 3+ | 3 | 3+ | 3 | 3+ | 4 | 4 |
| 19 | <i>TcLr13</i> | 3 | 3 | 3 | 3 | 4 | 3 | 3+ | 3 | 3 | 3 | 4 | 2 | 3 | 2 | 3 | 3+ | 2 | 2 |
| 20 | <i>TcLr14b</i> | 4 | 4 | 4 | 4 | 4 | 4 | 3+ | 4 | 4 | 4 | 4 | 3 | 4 | 4 | 3 | 4 | 4 | 4 |
| 21 | <i>TcLr15</i> | 4 | 1 | 1 | 1 | 3+ | 4 | 1 | 1 | 1 | 1 | 4 | 1 | 3 | 1 | 1 | 1 | 2 | 1, 2 |
| 22 | <i>TcLr19</i> | 0 | ; | 0 | 0 | 0 | 0 | 0 | ; | 0 | 0,; | 0 | ; | 0 | ; | 0 | 0 | 0 | 0 |
| 23 | <i>TcLr21</i> | 2 | 4 | 3 | 3 | 4 | 3 | 3 | ;;1 | 4 | 3+ | 3 | 3 | 3 | 4 | 3 | 2 | 3+ | 2+ |
| 24 | <i>TcLr20</i> | 1 | 1, 2 | 1 | 1, 2 | 1 | 1 | 1, 2 | ; | 1 | 1 | 1, 2 | 3 | 0 | 3+ | 1 | 3+ | 2+ | 1 |
| 25 | <i>TcLr23</i> | 1, 2 | 1 | 3 | 1 | 1 | 3 | 3 | 2+ | 3 | 3 | 1 | 1 | 1 | 1 | 1 | 3+ | 1 | 1 |
| 26 | <i>TcLr28</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | ; | 0 | ; | 0 | 0,; | 0 | 0 | ; | ; |
| 27 | <i>TcLr29</i> | 1 | 1 | 1 | 2 | 3 | 1 | 2 | 2 | 2 | 1 | 2 | ; | 2 | 3 | 3 | 1 | 3 | 2 |
| 28 | <i>TcLr33</i> | 3+ | 3+ | 3+ | 3+ | 3+ | 4 | 3+ | 4 | 3+ | 4 | 4 | 4 | 3 | 3+ | 3+ | 3+ | 3+ | 3 |
| 29 | <i>TcLr36</i> | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 3 | 1 | 2 | 1 | 1 | 1 | 3 | 1 | 1 |
| 30 | <i>TcLr39</i> | 3 | 2+ | 2+ | 2 | 3+ | 3 | 1 | 2 | 1 | 4 | 4 | 3 | 3 | 2 | 2 | 4 | 1 | 1 |
| 31 | <i>TcLr42</i> | 3 | 3 | 3 | 2 | 1 | 2 | 3 | 3+ | 3 | 1 | 1 | 2 | 2 | 2 | 3 | 1 | 3+ | 3 |
| 32 | <i>TcLr44</i> | 4 | 4 | 4 | 3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 3+ | 3 | 4 |
| 33 | <i>TcLr45</i> | 1 | 1 | 1 | ;;1 | 1 | 1 | 1 | 3 | ;;1 | 1 | 1 | 1 | 1 | 1 | 1 | 3+ | 1 | 1 |
| 34 | <i>TcLr47</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | ; | 0 | ; | ; | ; | ; | ; | 0 | ; |
| 35 | <i>TcLr51</i> | ; | ; | ; | 0 | ; | ; | ; | ; | ; | 1 | 1 | ; | 1 | 1 | ; | 1 | 1 | ; |
| 36 | <i>TcLr53</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | ; | 0 | ; | ; | ; | ; | ; | ; | 0,; |

^aAccording to the 0–4 Stakman scale modified by Roelfs et al. (1992); 0 – no flecks or uredinia, ; – hypersensitive flecks, 1 – small uredinia with necrosis, 2 – small uredinia with chlorosis, 3 – moderate size uredinia, 4 – large uredinia, + indicates slightly larger uredinia, C – more chlorosis than normal for the infection type, X – random distribution of variable-sized uredinia

<https://doi.org/10.17221/6/2021-CJGPB>

Table 3. Seedling infection types and the presence or absence of the *Lr* genes in 37 Chinese wheat cultivars based on the gene postulation using 18 *Puccinia triticina* (Pt) races and molecular markers

| No. ^b | <i>Lr</i> gene based on gene postulation | <i>Lr</i> gene based on gene marker detection | <i>Lr</i> gene(s) | Infection types ^a to <i>Pt</i> races | | | | | | | | | | | | | | | | | | | | | |
|------------------|--|---|---|---|----|------|------|------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| | | | | PH | TH | PH | JT | JS | PH | JS | PH | TH | PH | JT | JS | PH | TH | PH | JT | JS | PH | TH | PH | JT | JS |
| 1 | + | none | unknown APR genes | GQ | JT | PH | JT | JS | PH | JS | PH | TH | PH | JT | JS | PH | TH | PH | JT | JS | PH | TH | PH | JT | JS |
| | | | | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 2 | <i>Lr26, Lr13</i> , + | <i>Lr26</i> | <i>Lr26</i> ^c , <i>Lr13</i> ^d , + | 4 | 3 | 3+ | 3 | 4 | 3+ | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 |
| 3 | + | none | unknown seedling resistance gene | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 4 | <i>Lr1, Lr26</i> , + | <i>Lr1, Lr26</i> | <i>Lr1</i> ^c , <i>Lr26</i> ^c , + | 4 | 4 | 4 | 4 | 2 | 4 | 4 | 4 | 2 | 4 | 4 | 2 | 4 | 2 | 4 | 4 | 2 | 4 | 2 | 4 | 2 | 4 |
| 5 | + | none | unknown APR genes | 4 | 3 | 3 | 3 | 3+ | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 |
| 6 | <i>Lr1, Lr26</i> , + | <i>Lr1, Lr26</i> | <i>Lr1</i> ^c , <i>Lr26</i> ^c , + | 4 | 4 | 4 | 4 | 2 | 4 | 4 | 4 | 2 | 4 | 4 | 2 | 4 | 2 | 4 | 4 | 2 | 4 | 2 | 4 | 2 | 4 |
| 7 | <i>Lr1</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , + | 4 | 3+ | 4 | 4 | 2+ | 4 | 3 | 2+ | 2 | 4 | 3 | 2+ | 4 | 2 | 4 | 3 | 2 | 4 | 2 | 4 | 2 | 4 |
| 8 | <i>Lr1, Lr13</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , <i>Lr13</i> ^d , <i>Lr46</i> ^e , + | 3+ | 4 | 4 | 4 | 2 | 4 | 4 | 2 | 2 | 4 | 4 | 2 | 4 | 2 | 4 | 4 | 2 | 4 | 2 | 4 | 2 | 4 |
| 9 | + | <i>Lr46</i> | <i>Lr46</i> ^e , + | 3+ | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 10 | <i>Lr1, Lr13</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , <i>Lr13</i> ^d , + | 4 | 4 | 4 | 4 | 2 | 4 | 4 | 4 | 2c | 4 | 4 | 2 | 4 | 2 | 4 | 4 | 2 | 4 | 2 | 4 | 2 | 4 |
| 11 | + | none | unknown APR genes | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 12 | + | none | unknown APR genes | 4 | 3+ | 3 | 3 | 3+ | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 13 | <i>Lr1</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , + | 4 | 4 | 4 | 4 | 2+ | 4 | 3+ | 2 | 2 | 4 | 3 | 2 | 4 | 2 | 4 | 3 | 2 | 4 | 2 | 4 | 2 | 4 |
| 14 | + | none | unknown seedling resistance gene | 3 | 1 | 1, 2 | 1, 2 | 1, 2 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 |
| 15 | + | none | unknown APR genes | 4 | 4 | 4 | 3 | 3+ | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 16 | + | none | unknown APR genes | 4 | 4 | 4 | 3 | 3+ | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 17 | <i>Lr1</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , + | 3 | 4 | 2+ | 1 | 4 | 3 | 2+ | 1 | 1 | 3 | 3 | 2+ | 3 | 3 | 3 | 3 | 2+ | 3 | 3 | 3 | 3 | 3 |
| 18 | + | <i>Lr46</i> | <i>Lr46</i> ^e , + | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 19 | <i>Lr26</i> , + | <i>Lr26</i> | <i>Lr26</i> ^c , + | 4 | 3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 20 | + | none | unknown seedling resistance gene | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 21 | <i>Lr13, Lr18, Lr14a</i> , + | none | <i>Lr13</i> ^d , <i>Lr18</i> ^d , <i>Lr14a</i> ^d , + | 1 | 1 | 1 | 1 | 1, 2 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 | 1 | 4 |
| 22 | <i>Lr26</i> , + | <i>Lr26</i> | <i>Lr26</i> ^c , + | 3 | 3 | 3 | 3 | 3 | 4 | 3 | 4 | 4 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 |
| 23 | <i>Lr26</i> , + | <i>Lr26</i> | <i>Lr26</i> ^c , + | 3+ | 3 | 3 | 3 | 3+ | 4 | 3 | 4 | 4 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 |
| 24 | <i>Lr1</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , + | 3+ | 4 | 3 | 3 | 2+ | 4 | 4 | 2 | 1 | 2+ | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 |
| 25 | <i>Lr1</i> , + | <i>Lr1, Lr46</i> | <i>Lr1</i> ^c , <i>Lr46</i> ^e , + | 4 | 4 | 3 | 3 | 2 | 4 | 4 | 2 | 1 | 2 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 26 | + | <i>Lr34, Lr46</i> | <i>Lr34</i> ^e , <i>Lr46</i> ^e | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 27 | <i>Lr26</i> , + | <i>Lr26, Lr34</i> | <i>Lr26</i> ^c , <i>Lr34</i> ^e , + | 3+ | 4 | 3 | 3 | 3 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 28 | <i>Lr1</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , + | 4 | 3 | 3 | 3 | 2+ | 3 | 3 | 2+ | 2 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 |
| 29 | <i>Lr26</i> , + | <i>Lr26, Lr34</i> | <i>Lr26</i> ^c , <i>Lr34</i> ^e , + | 3 | 4 | 3 | 3 | 3 | 3 | 3 | 3 | 4 | 4 | 3 | 4 | 3 | 4 | 4 | 3 | 4 | 3 | 4 | 3 | 4 | 3 |
| 30 | <i>Lr26</i> , + | <i>Lr26</i> | <i>Lr26</i> ^c , + | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |

Table 3 to be continued

| No. ^b | Lr gene based on gene postulation | Lr gene based on gene marker detection | Lr gene(s) | Infection types ^a to <i>Pt</i> races | | | | | | | | | | | | | | | | | | | | | |
|------------------|-----------------------------------|--|------------------------------|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| | | | | PH | TH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH | PH |
| | | | | GQ | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT | JT |
| 31 | <i>Lr1</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , + | 4 | 3+ | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 32 | + | none | unknown APR gene | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 33 | <i>Lr26</i> , + | <i>Lr26</i> | <i>Lr26</i> ^c , + | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 34 | <i>Lr1</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , + | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 35 | <i>Lr1</i> , + | <i>Lr1</i> | <i>Lr1</i> ^c , + | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 36 | + | none | unknown APR genes | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 37 | + | none | unknown APR genes | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |
| 38 | – | none | CK | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 | 4 |

^aAccording to the 0–4 Stakman scale as modified by Roelfs et al. (1992); 0 – no flecks or uredinia, 1 – hypersensitive flecks, 2 – small uredinia with chlorosis, 3 – moderate size uredinia, 4 – large uredinia, + indicates slightly larger uredinia, C – more chlorosis than normal for the infection type, X – random distribution of variable-sized uredinia; ^bline numbers corresponding to those in Table 1; ^cpostulation of the *Lr* genes based on the gene postulation and molecular marker; ^dpostulation of the *Lr* genes based on the gene postulation; ^edetection of the *Lr* genes based on the molecular marker and adult plant resistance

RESULTS

***Lr* genes postulated and molecular marker detection.** The infection types of thirty-six differentials (Table 2), thirty-seven wheat cultivars and the susceptible control, Zhengzhou 5389 were evaluated with eighteen *Pt* races at the seedling stage in a greenhouse (Table 3). Zhengzhou 5389 showed IT 4 with all eighteen *Pt* races. The differential lines, containing *Lr9*, *Lr19*, *Lr24*, *Lr28*, *Lr47*, *Lr51*, and *Lr53* conferred a resistance response to all the *Pt* races, however, the differential lines carrying the *Lr* genes *2c*, *3*, *16*, *B*, *3bg*, *14b*, *33*, and *44* were susceptible to all the *Pt* races, hence, none of these genes could be postulated based the present response arrays. Twenty-one differential lines, *Lr1*, *Lr2a*, *Lr26*, *Lr3ka*, *Lr11*, *Lr17*, *Lr30*, *Lr10*, *Lr14a*, *Lr18*, *Lr2b*, *Lr13*, *Lr15*, *Lr20*, *Lr21*, *Lr23*, *Lr29*, *Lr36*, *Lr39*, *Lr42* and *Lr45* showed differential responses, and one or more of those genes could be postulated. The susceptible check, Zhengzhou 5389 was highly susceptible to all the *Pt* races (Table 3). Based on the IT arrays, *Lr* genes *1*, *13*, *14a*, *18* and *26*, either singly or in combination were postulated in twenty-two cultivars (Table 3). Fifteen cultivars were found with unknown resistance gene(s).

Lr26 was present in ten cultivars (Table 3). Five cultivars (Shaanken 6, Xiaoyan 166, Shaannong 138, Shaan 558 and Xiaoyan 228) contained only *Lr26* because they were only resistant to two *Lr26* avirulent races (TGGT and FGMT) and susceptible with the other sixteen *Pt* races. Two cultivars (E'mai 18 and Xiaoyan 22) contained *Lr26* combined with *Lr1* because they showed resistance to the *Lr26* and *Lr1* avirulent *Pt* races. Based on their resistant responses to other races, three cultivars (E'mai 17, Shaanmai 159 and Xi'nong 2611) had *Lr26* and other *Lr* genes in combination. Thirteen cultivars had *Lr1* based on the IT reactions (Table 3). Eight cultivars (Shaan 150, Xiaoyan 4, Shaannong 981, Xiaoyan 54, Xiaoyan 216, Xi'nong 9871, Xiaoyan 319 and Shaan 160) carried *Lr1* alone because, like the *Lr1* control, they were resistant to the same nine avirulent races (KHJS, KHHT, FHRT, FHJQ, FHTR, FHHTI, FHHTII, FHHT, and FGMT). Five lines contained *Lr1* combined with other *Lr* genes (*13*, *26* or *46*). All the cultivars with *Lr26* and *Lr1* were confirmed by the respective molecular markers for *Lr26* and *Lr1* (Table 3). Four cultivars contained *Lr13* because they had intermediate reactions to four *Lr13*-avirulent races (PHTTII, FHHTI, FHHT, and FGMT) (Table 3). Two cultivars, Gaoyou

<https://doi.org/10.17221/6/2021-CJGPB>

Table 4. Primer sequences and PCR amplification programmes for the different primer combinations

| <i>Lr</i> gene | Primer | | Cycle condition | Genetic distance (cM) | Size (bp) | Reference |
|----------------|-------------------|--------------------------|---|-----------------------|-----------|----------------------------|
| | Name | Sequence (5'–3') | | | | |
| <i>Lr1</i> | WR003F | GGGACAGAGACCTTGGTGGA | 94 °C 5 min; 35 cycles (94 °C 1 min; 55 °C 1 min; 72 °C 1 min); 72 °C 10 min; 10 °C forever | co-segregated | 745 | Qiu et al. (2007) |
| | WR003R | GACGATGATGATTTGCTGCTGG | | | | |
| <i>Lr9</i> | J13/1 | TCCTTTTATTCGACGCGCGG | 94 °C 6 min; 35 cycles (94 °C 1 min; 68.5 °C 1 min; 72 °C 2 min); 72 °C 10 min; 10 °C forever | co-segregated | 1 100 | Schachermayr et al. (1994) |
| | J13/2 | CCACACTACCCCAAAGAGACG | | | | |
| <i>Lr10</i> | FL2245 | GTGTAATGCATGCAGGTTCC | 94 °C 3 min; 35 cycles (94 °C 45 s; 60 °C 45 s; 72 °C 30 s); 72 °C 3 min; 10 °C forever | co-segregated | 282 | Schachermayr et al. (1997) |
| | <i>Lr10</i> -6/r2 | AGGTGTGAGTGAGTTATGTT | | | | |
| <i>Lr19</i> | SCS265-F | GGCGGATAAGCAGAGCAGAG | 94 °C 5 min; 35 cycles (94 °C 1 min; 65 °C 1 min; 72 °C 2 min); 72 °C 10 min; 10 °C forever | co-segregated | 525 | Gupta et al. (2006) |
| | SCS265-R | GGCGGATAAGTGGGTATG | | | | |
| <i>Lr19</i> | SCS253-F | GCTGGTTCCACAAAGCAAA | 94 °C 5 min; 35 cycles (94 °C 1 min; 60 °C 1 min; 72 °C 2 min); 72 °C 10 min; 10 °C forever | co-segregated | 736 | Gupta et al. (2006) |
| | SCS253-R | GGCTGGTTCCCTTAGATAGGTG | | | | |
| <i>Lr20</i> | STS638-L | ACAGCGATGAAGCAATGAAA | 94 °C 5 min; 35 cycles (94 °C 1 min; 60 °C 1 min; 72 °C 2 min); 72 °C 10 min; 10 °C forever | complete linkage | 500 | Neu et al. (2002) |
| | STS638-R | GTCCAGTTGGTTGATGGAAT | | | | |
| <i>Lr24</i> | J09/1 | TCTAGTCTGTACATGGGGGC | 94 °C 5 min; 35 cycles (94 °C 1 min; 60 °C 1 min; 72 °C 2 min); 72 °C 10 min; 10 °C forever | complete linkage | 315 | Schachermayr et al. (1995) |
| | J09/2 | TGGCACATGAACTCCATACG | | | | |
| <i>Lr26</i> | ω-secalinF | ACCTTCCTCATCTTTGTCCT | 94 °C 5 min; 35 cycles (94 °C 1 min; 65 °C 1 min; 72 °C 2 min); 72 °C 10 min; 10 °C forever | co-segregated | 1 100 | Chai et al. (2006) |
| | ω-secalinR | CCGATGCCATATACCACTACT | | | | |
| <i>Lr26</i> | O11B5 | GGTACCAACAACAACAACCC | 94 °C 5 min; 35 cycles (94 °C 1 min; 65 °C 1 min; 72 °C 2 min); 72 °C 10 min; 10 °C forever | co-segregated | 636 | Froidmont (1998) |
| | O11B3 | GTTGCTGCTGAGGTTGGTTC | | | | |
| <i>Lr34</i> | csLv34F | GTTGGTTAAAGACTGGTGATGG | 94 °C 5 min; 35 cycles (94 °C 1 min; 55 °C 1 min; 72 °C 2 min); 72 °C 10 min; 10 °C forever | 0.4 | 150 | Lagudah et al. (2006) |
| | csLv34R | TGCTTGCTATTTGCTGAATAGT | | | | |
| <i>Lr46</i> | csLV46G22-F | TCGACTTTGGGAATGGAGTTGC | 94 °C 5 min; 35 cycles (94 °C 1 min; 60 °C 1 min; 72 °C 2 min); 72 °C 10 min; 10 °C forever | complete linkage | 520 | Suenaga et al. (2001) |
| | csLV46G22-R | GGCGAAGATGCCATCATCCACCAG | | | | |

Table 5. Analysis of variance of the final leaf rust severities for 37 test lines and Saar and Zhengzhou 5389 controls tested over four growing seasons

| Source | df | MS | F | P value |
|--------------------------|-----|------------|--------------|---------|
| Cultivars | 38 | 4 710.609 | 3 740.173** | < 0.001 |
| Years | 3 | 8 496.879 | 6 746.431** | < 0.001 |
| Places | 1 | 30 240.692 | 2 4010.787** | < 0.001 |
| Replications | 1 | 4.858 | 3.857 | 0.051 |
| Cultivars × replications | 38 | 1.357 | 1.077 | 0.359 |
| Cultivars × environments | 230 | 554.123 | 439.967 ** | < 0.001 |
| Error | 234 | 1.259 | | |

$R^2 = 0.999$; **significance at level 0.01 probability; df – degree of freedom; MS – mean square

503 and Xi'an 93991, contained *Lr13* combined with *Lr1*. *Lr13* and *Lr26* were present in E'mai 17. Xi'nong 538 contained *Lr13*, *Lr18* and *Lr14a* in combination because they were resistant to races that were avirulent to the respective single gene controls. E'mai 12, Shaanmai 139 and Shaan 627 were resistant to some races and their resistance could not be attributed to any known *Lr* gene. E'mai 580, Xi'nong 291, Xi'nong 88, E'mai16, Xi'nong 2000, Xi'nong 223, Xi'nong 126, Xi'nong 3517, Xi'nong 6028, Xi'nong 889, Shaanmai 175 and Shaan 512 were susceptible to all the races (Table 3).

Nine molecular markers, eight STS and one SCAR closely linked to *Lr1*, *Lr9*, *Lr10*, *Lr19*, *Lr20*, *Lr24*, *Lr26*, *Lr34* and *Lr46* were used to genotype all the thirty-seven cultivars (Table 4). No line carried *Lr9*, *Lr10*, *Lr19*, *Lr20*, or *Lr24* based on the molecular marker

detection and gene postulation. The molecular marker tests predicted that *Lr46* was present in five cultivars Gaoyou 503, Xi'nong 88, Xi'nong 3517, Shaan 253 and Xi'nong 6028, And *Lr34* was detected in three cultivars (Xi'nong 6028, Shaanmai 159, and Xi'nong 2611) (Table 3). The cultivars containing *Lr34* and *Lr46* also showed symptoms of leaf tip necrosis (LTN) at the adult plant stage.

The combined results of the gene postulation and molecular marker detection indicated that seven *Lr* genes, 1, 26, 13, 14a, 18, 34, and 46, either singly or in combination were found in twenty-five lines, but twelve cultivars possibly contained unknown *Lr* gene(s) or lacked detectable *Lr* genes.

Adult plant resistance in field tests. The analysis of variance results of the FDS data (at $P = 0.05$) of the tested cultivar showed significant differences among

Table 6. Infection types (IT) in the seedling stage test with *Puccinia triticina* (Pt) races mixed Pt races and the mean final disease severity (FDS) in the field experiments with the same race in the 2014–2015, 2015–2016, 2016–2017 and 2017–2018 growing seasons for 37 wheat genotypes with the slow rusting resistance to leaf rust

| No. ^a | Cultivar (line) | Seedling ITs to race mixture ^a | FSD (%) | | | | | | |
|------------------|----------------------------|---|-----------|-------|-----------|-------|-----------|-------|-----------|
| | | | 2014–2015 | | 2015–2016 | | 2016–2017 | | 2017–2018 |
| | | | Baoding | Henan | Baoding | Henan | Baoding | Henan | |
| 2 | E'mai 17 | 4 | 2.5 | 15 | 2.5 | 20 | 5 | 15 | 5 |
| 3 | E'mai 12 | 4 | 2.5 | 17.5 | 2.5 | 20 | 1 | 15 | 35 |
| 16 | Xi'nong 126 | 4 | 12.5 | 25 | 12.5 | 15 | 5 | 1 | 15 |
| 17 | Shaannong 981 | 4 | 10 | 35 | 10 | 7.5 | 5 | 5 | 15 |
| 27 | Shaanmai 1591 | 4 | 20 | 7.5 | 20 | 15 | 1 | 20 | 10 |
| 36 | Shaanmai 175 | 4 | 10 | 15 | 10 | 10 | 7.5 | 2.5 | 8.75 |
| | Saar ^b | 4 | 2.5 | 1 | 5 | 1.75 | 5 | 1 | 1 |
| | Zhengzhou5389 ^c | 4 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| LSD | ($P = 0.05$) | | | | | | 22.17 | | |

^aLine numbers corresponding to those in Table 1; ^bslow rusting check; ^csusceptible check; LSD – least significance difference

<https://doi.org/10.17221/6/2021-CJGPB>

the cultivar-environment interaction, cultivars, and environments, but there were not significant differences between the cultivar-replication interaction (Table 5). In all the cropping seasons, the mean value of the FDS of the susceptible check Zhengzhou 5389 and the slow rusting check Saar was 100% and 2.5%, respectively, indicating an adequate level of disease. Six cultivars exhibiting high ITs to the *Pt* races mixture in the seedling tests showed consistent slow leaf rusting resistance in the field (Table 6). Among those cultivars, Shaanmai 159 carried *Lr34* and none of them carried *Lr46*.

DISCUSSION

The postulation of resistance genes based on the response to multiple races in seedling tests is a quick and traditional method of gene identification across diverse genetic backgrounds (Mebrate et al. 2008); and can be significantly strengthened by molecular marker genotyping and pedigree analysis. *Lr26* was derived from rye (*Secale cereal* L.) and located on the 1BR/1RS chromosome of wheat. *Lr26* is widely present in Chinese wheat lines in a high frequency through the introduction of wheat germplasm such as Lovrin 13, Lovrin 10, Predgoraia 2, Kavkaz, and Neuzucht (Zhuang 2003). According to Zhou et al. (2004), the frequency of the 1BR/1RS translocation line in the northern winter wheat area was 59%, and its frequency in the Huanghuai winter wheat area was 42%. In the present study, ten cultivars contained *Lr26* (Table 3). According to the pedigree analysis, *Lr26* in Xiaoyan 22 might be derived from Fengchan 3. Because Fengchan 3 carried *Lr26* (Li & Yan 1985; Zheng 2019).

The *Lr1* gene is widely distributed in various regions of the world such as Australia (McIntosh 1992), America (Kolmer et al. 2009) and Europe (Urbanovich et al. 2006). In China, *Lr1* was reported in more than 500 wheat varieties (Liu et al. 2014). Singh et al. (2000) also reported a high frequency of *Lr1* in Chinese cultivars and lines. In the present study, thirteen cultivars contained *Lr1*. The distribution frequencies of *Lr1* in Chinese cultivars are from four founder wheat parents, viz. the Nanda 2419, Funo, Yanda 1817 and Bima 4 derivatives (Liu et al. 2014). *Lr1* in Xiaoyan 216, Xi'nong 9871 and Xiaoyan 22 might be derived from Xiaoyan 22. Ren et al. (2012) reported that Xiaoyan 54, Xi'nong 9871, and Xiaoyan 22 contained *Lr1*. *Lr34* and *Lr46* are currently important leaf rust resistance genes in

China and are the most widely assessed slow rusting leaf rust resistance genes (Yuan & Chen 2011). *Lr34* and *Lr46* might not be highly effective when they are found alone, but they contribute higher levels of resistance when used in combination with other resistance genes (Sui et al. 2016).

In the present study, six cultivars were identified to carry slow rusting leaf rust resistance genes (Tables 3 and 6). E'mai 17, E'mai 12, Xi'nong 126, Shaannong 981, Shaanmai 1591 and Shaanmai 175 had other known *Lr* genes or unknown *Lr* genes. *Lr34*, *Lr46*, *Lr67* and *Lr68* are APR *Lr* genes, the molecular markers co-segregated with or closely linked to these were used to test these cultivars (Wu et al. 2019). E'mai 17 carried *Lr26* and *Lr13*, and this cultivar was found to have slow rusting based on the FDS. *Lr26* had lost resistance, although originally *Lr13* was described as an APR gene, which can be detected at the seedling stage especially at high temperatures (Pretorius et al. 1984). If *Lr26* and *Lr13* are found in combination, the resistance is significant.

The identified seedling or slow rusting *Lr* genes in the present tested wheat cultivars may facilitate the breeding process of Chinese wheat cultivars and might contribute to reducing the leaf rust damage in China.

CONCLUSION

In the study, seven *Lr* genes, *1*, *13*, *18*, *14a*, *26*, *34* and *46* either singly or in combination were identified in twenty-five lines. *Lr9*, *Lr10*, *Lr19*, *Lr20* and *Lr24* were not identified in all the tested cultivars. The known *Lr* genes, *9*, *19*, *24*, *28*, *47*, *51* and *53* were effective at all the plant growing stages. Five and three cultivars possessed *Lr46* and *Lr34*, respectively. Six cultivars showed slow leaf rusting resistance in the field. The results of this study are useful to incorporate the resistance genes from the sources identified here into the Chinese facultative wheat genotypes to improve the genetic diversity of the cultivars.

REFERENCES

- Chai J.F., Zhou R.H., Jia J.Z., Liu X. (2006): Development and application of a new co-dominant PCR marker for detecting 1BL/1RS wheat-rye chromosome translocations. *Plant Breeding*, 125: 302–304.
- Dehne H.W., Oerke E.C. (1998): Impact of diseases and disease control on crop production. In: Hutson D.H.,

- Miyamoto J. (eds.): Fungicidal Activity: Chemical and Biological Approaches to Plant Protection. Chichester, Wiley: 1–21.
- Dong J.G. (2001): Agricultural Plant Pathology. Beijing, China Agriculture Press.
- Dubin H.J., Torres E. (1981): Causes and consequences of the 1976–1977 wheat leaf rust epidemic in north west Mexico. *Annual Review of Phytopathology*, 19: 41–49.
- Flor H.H. (1955): Host-parasite interaction in flax rust-its genetics and other implications. *Phytopathology*, 45: 680–685.
- Froidmont D.D. (1998): A codominant marker for the 1BL/1RS wheat-rye translocation via multiplex PCR. *Journal of Cereal Science*, 27: 229–232.
- Fu D.L., Uauy C., Distelfeld A., Blechl A., Epstein L., Chen X., Sela H., Fahima T., Dubcovsky J. (2009): A Kinase-START gene confers temperature-dependent resistance to wheat stripe rust. *Science*, 323: 1357–1360.
- Gebrewahid T.W., Yao Z.J., Yan X.C., Gao P., Li Z.F. (2017): Identification of leaf rust resistance genes in Chinese common wheat cultivars. *Plant Disease*, 101: 1729–1737.
- Gupta S.K., Charpe A., Prabhu K.V., Haque Q.M.R. (2006): Identification and validation of molecular markers linked to the leaf rust resistance gene *Lr19* in wheat. *Theoretical and Applied Genetics*, 113: 1027–1036.
- Helguera M., Khan I.A., Dubcovsky J. (2000): Development of PCR markers for the wheat leaf rust resistance gene *Lr47*. *Theoretical and Applied Genetics*, 100: 1137–1143.
- Kolmer J.A., Long D.L., Hughes M.E. (2009): Physiologic specialization of *Puccinia triticina* on wheat in the United States in 2007. *Plant Disease*, 93: 538–544.
- Lagudah E.S., McFadden H., Singh R.P., Huerta-Espino J., Bariana H.S., Spielmeier W. (2006): Molecular genetic characterization of the *Lr34/Yr18* slow rusting resistance gene region in wheat. *Theoretical and Applied Genetics*, 114: 21–30.
- Li Z., Yan Z.L. (1985): Selection and utilization of dwarf wheat Xiaoyan 392 in breeding. *Shaanxi Agricultural Science*, 5: 27–28.
- Li Z.F., Xia X.C., He Z.H., Li X., Zhang L.J., Wang H.Y., Meng Q.F., Yang W.X., Li G.Q., Liu D.Q. (2010): Seedling and slow rusting resistance to leaf rust in Chinese wheat cultivars. *Plant Disease*, 94: 45–53.
- Li Z.F., Lan C.X., He Z.H., Singh R.P., Rosewarne G.M., Chen X.M., Xia X.C. (2014): Overview and application of QTL for adult plant resistance to leaf rust and powdery mildew in wheat. *Crop Science*, 54: 1907–1925.
- Lillemo M., Asalf B., Singh R.P., Huerta-Espino J., Chen X.M., He Z.H., Bjørnstad Å. (2008): The adult plant rust resistance loci *Lr34/Yr18* and *Lr46/Yr29* are important determinants of partial resistance to powdery mildew in bread wheat line Saar. *Theoretical and Applied Genetics*, 116: 1155–1166.
- Liu X.C., Lai Y.P., Liu X.J., Li J., Yuan J.E., Wang H.S., Feng Z.Y., Yang W.Y. (2014): Distribution and selective effects of *Lr1* gene in the wheat varieties derived from four founder breeding parents in China. *Journal of Triticeae Crops*, 34: 597–602.
- Long D.L., Kolmer J.A. (1989): A north American system of nomenclature for *Puccinia triticina*. *Phytopathology*, 79: 525–529.
- McIntosh R.A. (1992): Pre-emptive breeding to control wheat rusts. *Euphytica*, 63: 1–2.
- McIntosh R.A., Wellings C.R., Park R.F. (1995): Wheat Rusts: An Atlas of Resistance Genes. East Melbourne, CSIRO Publications.
- Mebrate S.A., Dehne H.W., Pillen K., Oerke E.C. (2008): Postulation of seedling leaf rust resistance genes in selected Ethiopian and German bread wheat cultivars. *Crop Science*, 48: 507–516.
- Neu C., Stein N., Keller B. (2002): Genetic mapping of the *Lr20-Pm1* resistance locus reveals suppressed recombination on chromosome arm 7AL in hexaploid wheat. *Genome*, 45: 737–744.
- Peng H., Lv G.Q., Wang J.R. (2016): Analysis on the characteristics and causes of main diseases of wheat in Henan Province in 2015. *China Plant Protection*, 36: 29–33.
- Pretorius Z.A., Wilcoxson R.D., Long D.L., Schafer J.F. (1984): Detecting wheat leaf rust resistance gene *Lr13* in seedlings. *Plant Disease*, 68: 585–586.
- Qiu J.W., Schvrch A.C., Yahiaoui N., Dong L.L., Fan H.J., Zhang Z.J., Keller B., Ling H.Q. (2007): Physical mapping and identification of a candidate for the leaf rust resistance gene *Lr1* of wheat. *Theoretical and Applied Genetics*, 115: 159–168.
- Peterson R.F., Campbell A.B., Hannah A.E. (1948): A diagrammatic scale for estimating rust intensity on leaves and stems of cereals. *Canadian Journal of Research*, 26: 496–500.
- Ren X.L., Liu T.G., Liu B., Gao L., Chen W.Q. (2012): Multiplex PCR assay for detection of wheat leaf rust resistance genes *Lr9-Lr29* and *Lr19-Lr20* in 116 Chinese wheat cultivars (lines). *Plant Protection*, 38: 29–36.
- Roelfs A.P., Singh R.P., Saari E.E. (1992): Rust Diseases of Wheat: Concepts and Methods of Disease Management. Mexico City, CIMMYT.
- Schachermayr G., Siedler H., Gale M.D., Winzeler H., Winzeler M., Keller B. (1994): Identification and localization of molecular markers linked to the *Lr9* leaf rust resistance gene of wheat. *Theoretical and Applied Genetics*, 88: 110–115.
- Schachermayr G.M., Messmer M.M., Feuillet C., Winzeler H., Winzeler M., Keller B. (1995): Identification of

<https://doi.org/10.17221/6/2021-CJGPB>

- molecular markers linked to the *Agropyron elongatum*-derived leaf rust resistance gene *Lr24* in wheat. Theoretical and Applied Genetics, 90: 982–990.
- Schachermayr G., Feuillet C., Keller B. (1997): Molecular markers for the detection of the wheat leaf rust resistance gene *Lr10* in diverse genetic backgrounds. Molecular Breeding, 3: 65–74.
- Sharp P.J., Kreis M., Shewry P.R. (1988): Location of β -amylase sequence in wheat and its relatives. Theoretical and Applied Genetics, 75: 286–290.
- Singh R.P., Chen W.Q., He Z.H. (1999): Leaf rust resistance of spring, facultative, and winter wheat cultivars from China. Plant Disease, 83: 644–651.
- Singh R.P., Huerta-Espino J., Rajaram S. (2000): Achieving near-immunity to leaf and stripe rusts in wheat by combining slow rusting resistance genes. Acta Phytologica Sinica, 35: 133–139.
- Suenaga K., Singh R.P., Manilal H.M. (2001): Tagging of slow rusting genes for leaf rust, *Lr34* and *Lr46*, using microsatellite markers in wheat. JIRCAS Research Highlights, 2021: 8–9.
- Sui J.S., Wang H.L., Xin Z.H., Wang W., Chen T.Q., He Q.C. (2016): Detection of molecular markers of leaf rust resistance genes *Lr26*, *Lr34* and *Lr38* in 122 wheat varieties (lines). Seed, 5: 35–41.
- Urbanovich O.Y., Malyshev S.V., Dolmatovich T.V., Kartel N.A. (2006): Identification of leaf rust resistance genes in wheat (*Triticum aestivum* L.) cultivars using molecular markers. Russian Journal of Genetics, 42: 546–554.
- Wu H., Kang Z.H., Li X., Li Y.Y., Li Y., Wang S., Liu D.Q. (2019): Identification of wheat leaf rust resistance genes in Chinese wheat cultivars and the improved germplasms. Plant Disease, 101: 1729–1737.
- Yuan J.H., Chen W.Q. (2011): Evaluation of the effectiveness of main wheat leaf rust resistance genes in China. Journal of Triticeae Crops, 31: 994–999.
- Yuan J.H., Liu T.G., Chen W.Q. (1992): Postulation of leaf rust resistance genes in 47 new wheat cultivars (lines) at seeding stage. Scientia Agriculture Sinica, 22: 369–375.
- Zheng R.M. (2019): Molecular Detection and Genetic Derivation of 262 Chinese Wheat Microcore Germplasm against Leaf Rust. Yangling, Northwest A & F University.
- Zhou H.X., Xia X.C., He Z.H., Li X., Wang C.F., Li Z.F., Liu D.Q. (2013): Molecular mapping of leaf rust resistance gene *LrNJ97* in Chinese wheat line Neijiang 977671. Theoretical and Applied Genetics, 126: 2141–2147.
- Zhou Y., He Z.H., Zhang G.S., Xia L.Q., Chen X.M., Gao Y.C., Jing Z.B., Yu G.J. (2004): Utilization of 1BL/1RS translocation in wheat breeding in China. Acta Agonomica Sinica, 30: 531–535.
- Zhuang L.J., Li Z.F., Lillemo M., Xia X.C., Liu D.Q., Yang W.X., Luo J.C., Wang H.Y. (2009): QTL mapping for adult-plant resistance to leaf rust in CIMMYT wheat cultivar Saar. Scientia Agriculture Sinica, 42: 388–397.
- Zhuang Q.S. (2003): Chinese Wheat Improvement and Pedigree Analysis. Beijing, China Agriculture Press.

Received: January 24, 2021

Accepted: April 12, 2021

Published online: May 6, 2021